

Wzory na pochodne:

1. $(C)' = 0$
2. $(x^n)' = nx^{n-1}$
3. $(x)' = 1$
4. $\left(\frac{1}{x}\right)' = -\frac{1}{x^2}$
5. $(\sqrt{x})' = \frac{1}{2\sqrt{x}}$
6. $(a^x)' = a^x \ln a$
7. $(e^x)' = e^x$
8. $(\log_a x)' = \frac{1}{x \ln a}$
9. $(\ln x)' = \frac{1}{x}$
10. $(\sin x)' = \cos x$
11. $(\cos x)' = -\sin x$
12. $(\operatorname{tg} x)' = \frac{1}{\cos^2 x}$
13. $(\operatorname{ctg} x)' = -\frac{1}{\sin^2 x}$
14. $(\arcsin x)' = \frac{1}{\sqrt{1-x^2}}$
15. $(\arccos x)' = -\frac{1}{\sqrt{1-x^2}}$
16. $(\operatorname{arctg} x)' = \frac{1}{x^2+1}$
17. $(\operatorname{arcctg} x)' = -\frac{1}{x^2+1}$

Właściwości pochodnych:

1. $[f(x) + g(x)]' = f'(x) + g'(x)$
2. $[f(x) - g(x)]' = f'(x) - g'(x)$
3. $[a \cdot f(x)]' = a \cdot f'(x)$
4. $[f(x) \cdot g(x)]' = f'(x)g(x) + f(x)g'(x)$
5. $\left[\frac{f(x)}{g(x)}\right]' = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$

Wzory przydatne w liczeniu pochodnych:

$$\sqrt[b]{x^a} = x^{\frac{a}{b}}$$

$$\frac{1}{x^a} = x^{-a}$$